

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 25/30, 57/20	A1	(11) International Publication Number: WO 99/17608 (43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: PCT/GB98/02951 (22) International Filing Date: 1 October 1998 (01.10.98) (30) Priority Data: 9720891.2 2 October 1997 (02.10.97) GB (71) Applicant (for all designated States except US): IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chemical House, Millbank, London SW1P 3JF (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): AUDA, Mahroussa [BE/BE]; Stijn Streuvelsstraat 6, B-9051 Siont Denijs Westrem (BE). HARTMANN, Frank, Dirk, Jozef [BE/BE]; Mellesteenweg 10, B-9230 Wetteren (BE). REEKMANS, Steven, Irene, Jozef [BE/BE]; Dejonker Straat 51, B-1060 Brussels (BE). (74) Agents: ROBERTS, Jonathan, Winstanley et al.; ICI Group Intellectual Property, The Heath, P.O. Box 11, Runcorn, Cheshire WA7 4QE (GB).		(81) Designated States: AU, BR, CA, CN, CZ, HU, ID, JP, KR, MX, NZ, PL, RU, SG, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: AGROCHEMICAL COMPOSITIONS (57) Abstract Adjuvants for aqueous agrochemical compositions incorporating hydrocarbyl polysaccharides can be formulated with polyalkylene glycols, particularly polyethylene glycols, without a major loss of effectiveness in the total formulation. The compositions can include other water soluble or dispersible non-ionic surfactants. The agrochemical compositions can be used to kill or inhibit vegetation where the agrochemical includes a growth regulator and/or herbicides, particularly water soluble herbicides, such as a glyphosate type herbicides; or to kill or inhibit plant pests where the agrochemical includes a pesticide, particularly an insecticide, fungicide and/or acaricide.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Agrochemical compositions

This invention relates to agrochemical compositions and to adjuvant compositions for agrochemical compositions including alkyl saccharide surfactants. It further relates to the use of such adjuvant compositions in agrochemical compositions especially compositions containing growth regulators, herbicides, insecticides, fungicides or acaricides, particularly for spray application.

Hydrocarbyl saccharide surfactants are known materials in which a hydrocarbyl group, typically an alkyl group, is attached to a saccharide residue by an ether bond. The saccharide residue often either is or contains more than one saccharide unit and this class of surfactant is often referred to as being hydrocarbyl (alkyl) polysaccharides. As the number of saccharide repeat residues is not usually very large, often containing low levels of chains having more than 4 saccharide units and typically having an average number of between 1 and 2, describing such materials as polysaccharides is something of a misnomer, although the terminology survives widely in the art. We use the term hydrocarbyl (or alkyl) saccharide to refer to this class of surfactant.

Hydrocarbyl saccharide surfactants are effective adjuvants for agrochemical, particularly growth regulator and/or pesticide active compounds, particularly for glyphosate type herbicides and their use in this way is described in EP 0220902 A. Formulations additionally including other, particularly non-ionic, surfactants are described in EP 0671967 A (WO 94/12259 A) and such other surfactants can contribute to the adjuvancy in the overall system. Typically hydrocarbyl saccharides are used as adjuvants (particularly in end use spray formulations) in solution in water. The present invention is based on our discovery that by using relatively low molecular weight polyethylene glycols (PEGs) as a component of the adjuvant system, the overall amount of hydrocarbyl saccharide used can be reduced without loss of adjuvant activity. This is very surprising because PEGs are not in themselves surfactants and are not recognised as adjuvants and would thus be expected simply to dilute the effectiveness of the hydrocarbyl saccharides.

Accordingly, the present invention provides a composition which comprises

- i at least one hydrocarbyl polysaccharide;
- ii at least one polyalkylene, particularly at least one polyethylene, glycol; and
- iii water.

The invention further includes a water dispersible and/or soluble agrochemical composition comprising:

- i at least one hydrocarbyl polysaccharide;

- 2 -

- ii at least one polyalkylene, particularly at least one polyethylene, glycol;
- iii water; and
- iv at least one agrochemical.

In this aspect of the invention, the agrochemical is typically one or more plant growth regulators, herbicides, and/or pesticides, for example insecticides, fungicides or acaricides. The invention is particularly applicable to water soluble agrochemicals and especially to such agrochemicals that form aqueous solutions of high ionic strength. Notable examples of such agrochemicals are the glyphosate type of herbicide. The invention accordingly includes an agrochemical composition which is dispersible in water and is itself a stable homogeneous aqueous solution comprising:

- i at least one hydrocarbyl polysaccharide;
- ii at least one polyalkylene, particularly at least one polyethylene, glycol;
- iii water; and
- v at least one glyphosate type herbicide.

The composition can include other surfactants, particularly non-ionic surfactants so that the invention further includes a composition which comprises

- i at least one hydrocarbyl polysaccharide;
- ii at least one polyalkylene, particularly at least one polyethylene, glycol; and
- iii water, and
- iv at least one other water soluble or water dispersible non-ionic surfactant.

The main reason for including such other water soluble or water dispersible non-ionic surfactants is for their contribution to overall adjuvancy in agrochemical compositions. The invention thus includes a water dispersible and/or soluble agrochemical composition comprising:

- i at least one hydrocarbyl polysaccharide;
- ii at least one polyalkylene, particularly at least one polyethylene, glycol;
- iii water;
- iv at least one other water soluble or water dispersible non-ionic surfactant; and
- v at least one agrochemical.

This aspect of the invention also applies to water soluble agrochemicals that form aqueous solutions of high ionic strength, particularly glyphosate type herbicides and the invention accordingly further includes an agrochemical composition which is dispersible in water and is itself a stable homogeneous aqueous solution comprising:

- i at least one hydrocarbyl polysaccharide;
- ii at least one polyalkylene, particularly at least one polyethylene, glycol;
- iii water;
- iv at least one other water soluble or water dispersible non-ionic surfactant ;and

- 3 -

- v at least one glyphosate type herbicide.

The invention includes methods of use of the adjuvant and agrochemical compositions of the invention specifically a method of treating vegetation by applying to plants and/or soil near the plants an agrochemical formulation according to the invention. More particularly, the invention includes the use of agrochemical compositions including adjuvant compositions of this invention in the control of pests and of plant growth. Accordingly, the invention includes:

- (i) a method of killing or inhibiting vegetation by applying to the vegetation or the soil near or surrounding the vegetation an agrochemical composition of the invention where the agrochemical is or includes one or more growth regulators and/or herbicides, particularly including at least one glyphosate type herbicide; and/or
- (ii) a method of killing or inhibiting plant pests by applying to the vegetation or the soil surrounding the vegetation an agrochemical composition of the invention where the agrochemical is or includes one or more pesticides, for example insecticides, fungicides or acaricides.

The hydrocarbyl saccharide is typically at least one compound of the formula:



R is a hydrocarbyl group;

G is a saccharide residue; and


a is an average value which is at least 1.

R can be a substituted or unsubstituted hydrocarbyl group, such as a cycloalkyl, aryl, alkaryl, aralkyl or alkenyl or, and preferably, an alkyl, group, and suitably contains from 4 to 30, more usually up to 24, particularly from 6 to 18, especially 8 to 14, carbon atoms. Particularly as the compounds providing the group R are typically derived from natural products or distillation cuts, R can be a mixture of alkyl groups as described above and which may contain, on average, for example 8 to 14 carbon atoms. Deliberate mixtures of alkyl groups can also be used.

The saccharide residue G can particularly be a glycoside and especially a glucoside residue. Thus it may be derived from one more of fructose, glucose, mannose, galactose, telose, gulose, allose, altrose, idose, arabinose, xylose, lyxose and ribose or from mixtures thereof. The group G is particularly conveniently derived from glucose units and the saccharide is then a glucoside. If derived from sucrose the groups will comprise fructose and glucose residues.

- 4 -

The value of a is the degree of polymerisation of the hydrocarbyl (poly)saccharide and as it is an average, it may and often will be non-integral. It is usually greater than 1, typically at least 1.1, desirably at least 1.2, particularly at least 1.3, and is typically not greater than 8, more usually not greater than 4, for example not greater than 2. In particular, when the saccharide is an alkyl glucoside, a is conveniently between 1 and 2. We have obtained useful results using alkyl glucosides of the general formula:



R' is a C_8 to C_{14} alkyl group; and

a' is between 1 and 2, preferably from about 1.3 to about 1.9.

In particularly useful alkyl glucosides R' is a C_9 to C_{13} , particularly about C_{10} , alkyl group, or a mixture of alkyl groups having an average carbon chain length in this range and in an especially useful alkyl glucoside R' is about C_{10} alkyl from mixed C_8 to C_{11} alkyl groups and a' is about 1.3 to about 1.9.

Hydrocarbyl glycosides are commercially available materials designated for example as Atplus 452, Atplus 508 from ICI Surfactants or as sold under the trade name Triton BG 10 by Rohm & Haas.

The polyalkylene glycols used are desirably polyethylene glycols (PEGs), polypropylene glycols or co-poly(ethylene/propylene) glycols. Generally the glycols have a molecular weight of from about 200 and usually not more than about 2000, more usually from about 250 to about 1000. When polyethylene glycol(s) are used they typically have molecular weights of from about 200 up to about 2000, more usually from about 250 to about 1000, and particularly not more than about 800. These molecular weight ranges correspond to numbers of repeat units (n) in the PEG chain of about 4 to about 45, more usually about 5 to about 22 and preferably not more than about 18. Particularly useful PEGs are those with molecular weights in the range 250 to 600 (n = about 5 to about 16), especially about 300 to about 400 (n = about 6.5 to about 8.5). For ease of handling it is desirable to use liquid PEGs and this corresponds (depending on ambient temperature and method of manufacture) to PEGs having molecular weights not more than about 800. Polypropylene glycols can have similar molecular weights, but more usually will not have molecular weights as high as those possible with polyethylene glycols. Typically molecular weights will be in the range 200 to 600, especially about 250 to about 400.


The weight ratio of hydrocarbyl saccharide to polyalkylene glycol, particularly PEG, used is desirably in the range of about 20:1 to 1:2. Proportions of polyalkylene glycol, particularly PEG,

- 5 -

lower than about 20:1 are not significantly more economic than the hydrocarbyl saccharide itself. Increasing the proportion of polyalkylene glycol, particularly PEG, above about 1:1 results in diminished performance at constant adjuvant additive use levels. The amount of the adjuvant composition could be increased relative to the agrochemically active components of an end use formulation to restore the performance, but at the cost of reducing or eliminating the economic benefit. Preferred ranges for the ratio of hydrocarbyl saccharide to polyalkylene glycol, particularly PEG, are from about 10:1 to about 1.5:1, particularly from about 5:1 to about 2:1.

Particularly when the agrochemically active components are or include highly ionic components, particularly when glyphosate type herbicides are used, and especially where the concentration of the solutes is high e.g. as in a so-called adjuvant built in concentrate (intended for dilution prior to use) the proportion of polyalkylene glycol, particularly PEG, may be limited if the molecular weight is above about 400, particularly above about 500 in order to maintain compatibility in the concentrate composition. Desirably, in such formulations, the polyalkylene glycol, particularly PEG, has a molecular weight of from about 200 to about 500, particularly about 250 to about 400.

The formulation can include other surfactants and in particular non-ionic surfactants of the formulae:



R^1 is a C_8 to C_{18} alkyl, (C_6 to C_{12} alkyl) alkyl phenyl, a sorbitan or a C_{10} to C_{22} fatty acyl group;

R^2 is hydrogen or an alkyl or carboxyalkyl group or a salt thereof, or a C_{10} to C_{22} fatty acyl group;

X is a polyalkylene oxide group having an average of 2 to 40 alkylene oxide groups or mixed polyalkylene oxide groups, and

R^3 is a C_{10} to C_{22} alkyl or alkenyl group.

Preferred non-ionic surfactants of these formulae include those in which

R^1 is a C_{12} to C_{15} alkyl group, or a sorbitan group,

R^2 is hydrogen, a C_1 to C_4 alkyl, or a carboxymethyl group or a salt thereof, or a C_{10} to C_{22} fatty acyl group;

X is a polyalkylene oxide, particularly polyethylene or polypropylene oxide, group having an average of 3 to 30 alkylene oxide residues or mixed polyalkylene oxide, particularly mixed polyethylene or polypropylene oxide, group, containing 2 to 25, and preferably 3 to 20 alkylene oxide residues; and

- 6 -

R^3 is a C_{16} to C_{18} alkyl or alkenyl group.

Examples of preferred non-ionic surfactants thus include poly(alkoxylated) C_8 to C_{24} , preferably C_{12} to C_{18} fatty acids and alcohols; sorbitan and sorbitol esters, e.g. sorbitan monolaurate and monooleate; poly(alkoxylated) derivatives of sorbitan and sorbitol esters; and poly(alkoxylated) fatty amines; and combinations of two or more such compounds. Among poly(alkoxylated) alcohols, polyethoxylates of alcohols with C_{12} to C_{18} straight chain alkyl and primary monobranched groups are particularly useful and among poly(alkoxylated) fatty amines, tallow amine ethoxylates are particularly useful, because they can be incorporated into concentrate compositions which are stable and homogeneous aqueous solutions. Ethoxylated sorbitan esters, particularly the oleate and stearate, e.g. the materials sold by ICI under the trade name Tween, are useful, but do not readily form stable homogeneous aqueous solution concentrate compositions and are thus useful primarily as additives when making up the final mix for spraying i.e. as so called "tank mix additives".



The amount of any other non-ionic surfactant used can vary widely up to about 1.5 times the weight amount of alkyl glycoside used. Typically, the amount, when used, will be from 5% to 120%, more usually from 5 to 100%, by weight of the alkyl glycoside. Generally the total of any other non-ionic surfactant and the polyalkylene glycol will not exceed 150% of the weight of the alkyl glycoside used.

In adjuvant compositions of the invention, the weight ranges of composition will usually be in the following ranges:

Component	Range			
	Broad (wt%)		Narrow (wt%)	
Hydrocarbyl (particularly alkyl) saccharide	25	70	35	70
PAG (particularly PEG)	2	35	5	30
Other non-ionic surfactant (when present)	2	35	5	30
Minor components	0	5	0	5
Water	15	70	20	60

The agrochemical used in this invention is typically one or more plant growth regulators, herbicides, and/or pesticides, for example insecticides, fungicides or acaricides. Particular agrochemicals which can be used in this invention include glyphosate type herbicides which are typically compounds of the formula:

- 7 -



R is halogen, -NHOH, -N(R¹)₂, -OR₂, -SR₂ or -OM,

where each R¹ is independently hydrogen, or an alkyl, hydroxyalkyl, or alkenyl group, preferably containing up to about 5 carbon atoms or a phenyl group;

each R² is independently hydrogen, or an alkyl, hydroxyalkyl, chloroalkyl, or alkoxy group, preferably containing up to about 5 carbon atoms, an alkylene amine group, preferably containing up to about 12 carbon atoms, or a phenyl or benzyl group;

M is hydrogen or an agriculturally acceptable salt forming moiety, such as alkali or alkaline earth metal, or a stannic, ammonium, organic ammonium, alkyl sulfoxonium, or alkyl phosphonium moiety or a combination thereof; and

Z is hydrogen, an organic moiety or an inorganic moiety.

Such compounds are described in US Patents 3799758, 4397676, 4140513, 4315765, 3868407, 4405531, 4481026, 4414158, 4120689, 4472189, 4341549 and 3948975 and, where Z is other than hydrogen, US Patents 3888915, 3933946, 4062699, 4119430, 4322239 and 4084954. In preferred compounds, Z is hydrogen or an organic substituent. Suitable organic substituents include methylene carboxylic; methylene phosphonic; methylene cyano; carboxyl, such as formyl, acetyl, benzoyl, perfluoroacyl and thiocarbonyl; ethylene, such as cyano, carbamoyl or carboxy substituted ethyl; and benzene sulphonyl substituents. Patents disclosing compounds where the nitrogen contains three organic substituents include US Patents 3455675, 3556762, 3853530, 3970695, 3988142, 3991095, 3996040, 4047927, 4180394, 4203756, 4261727 and 4312662. A preferred tertiary nitrogen substituted compound is N,N-bis(phosphonomethyl)glycine. Compounds where Z is hydrogen are particularly desirable when the phytoactivity desired is herbicidal activity.

Agriculturally acceptable salt forming moieties represented by M, as in OM, include alkali metals particularly sodium, potassium, or rubidium; alkaline earth metals particularly magnesium or calcium; ammonium and aliphatic ammonium, wherein the aliphatic is primary, secondary, tertiary or quaternary and preferably where the total number of carbon atoms is not more than about 12; phenylammonium; trialkylsulphonium, preferably where the total number of carbon atoms in the three alkyl substituents is not more than about 6, such as trimethylsulphonium, ethyl dimethylsulphonium, propyl dimethylsulphonium and the like; trialkylsulphoxonium, preferably where the total number of carbon atoms in the three alkyl substituents is not more than about 6, such as trimethylsulphoxonium, ethyl dimethylsulphoxonium, propyl dimethylsulphoxonium and the like;

- 8 -

tetraalkylphosphonium, such as tetramethylphosphonium, ethyl trimethylphosphonium, propyl trimethylphosphonium and the like.

In preferred formulations according to this invention, M is independently an agriculturally acceptable salt forming moieties e.g. as described above or hydrogen, particularly alkali metal, ammonium, monoalkyl ammonium or trialkylsulphonium. In particularly desirable formulations one M is an alkali metal, ammonium, monoalkyl ammonium, or trialkylsulphonium moiety, and the other two are hydrogen. Such particularly desirable compounds include *iso*-propylamine *N*-phosphonomethylglycine, trimethylsulphonium *N*-phosphonomethylglycine and sodium sesqui-*N*-phosphonomethylglycine. Combinations of two or more such compounds can be used in this invention.

Agrochemical compositions are often formulated as concentrates which are intended to be diluted with water immediately before use. The invention includes agrochemical compositions in the form of dilutable concentrates. As embodied in stable aqueous dilutable solutions containing water soluble agrochemicals, the invention relates to such concentrated compositions particularly having concentrations of the various components in the following ranges.

Component	Range (g.l ⁻¹)	
agrochemical	100	500
adjuvant **	30	500
minor components	0	50
water	remainder	

As embodied by aqueous compositions containing glyphosate type herbicides typical ranges are:

Component	Range (g.l ⁻¹)		Typical (g.l ⁻¹)
glyphosate herbicide expressed as:			
acid	100	500	330
salt used*	145	725	480
adjuvant **	30	500	240
minor components	0	50	as needed
water	remainder		

the ratio of glyphosate (as acid) : adjuvant being from 10:1 to 1:3

* calculated as for sulfosate (= glyphosate trimethylsulphonium salt)

** here adjuvant includes the hydrocarbyl saccharide, the polyalkylene glycol and any other non-ionic surfactant present.

The invention includes agrochemical concentrate compositions based on glyphosate type herbicides and this forms a specific aspect of the invention which accordingly provides: an

- 9 -

agrochemical concentrate composition which is dispersible in water and is itself a stable homogeneous aqueous solution comprising:

from 30 to 500 g.l⁻¹ of an adjuvant composition comprising

- a at least one hydrocarbyl polysaccharide;
- b at least one polyalkylene, particularly at least one polyethylene, glycol;
- c optionally at least one other water soluble or water dispersible non-ionic surfactant;
- and
- d water; and

from 100 to 500 g.l⁻¹ of at least one glyphosate type herbicide.

In comparison, the concentrations in the spray formulations will be much lower, typically of the order of 100 times e.g. 50 to 1000 times, more dilute. The relative concentrations will generally remain as indicated above. The particular concentration in any specific case will be determined by the user based on the desired dose of agrochemical to be applied and on the volume of spray to be used. The spray volume will depend on the spraying method and equipment used, but using current methods the spray volume will typically be from about 100 to 400 l.ha⁻¹.

The compositions of the invention may also incorporate other components such as antifoams, viscosity modifiers, stabilisers and antimicrobials.

When present the antifoam component is desirably a polysiloxane antifoam agent, e.g. of the

formula: $R^{10}_3SiO - [SiO R^{10} R^{11}]_n - Si R^{10}_3$ in which the groups R¹⁰ are each alkyl,

preferably C₁ to C₃ alkyl, especially methyl, groups, and one or more of the groups R¹¹ may be polyalkylene glycol residues and the others as defined for R¹⁰. The value of n is desirably such that the polysiloxanes have a molecular weight of from 1000 to 20000, preferably 5000 to 15000. Examples of suitable polysiloxane antifoams are disclosed in British Patents 1533610 and 1554736. The amount of antifoam used should be sufficient to secure an adequate anti-foaming effect in use and is normally 0.1 to 10% and preferably 0.2 to 6% by weight of the surfactants present in the formulation corresponding to about 0.01 to 5%, particularly 0.02 to 2%, by weight of an adjuvant or agrochemical concentrate and typically 0.0001 to 0.1%, preferably 0.001 to 0.05%, by weight of a spray formulation at end use dilution.

- 10 -

Other possible additives in the dispersible and/or soluble concentrate to which this invention relates include non-surfactant materials which are conventionally useful in surfactant formulations, such as viscosity modifiers, stabilisers, and anti-microbials. One class of known viscosity modifier materials of this type includes one or more commercially available water soluble or miscible materials such as gums, e.g. xanthan gums, and/or cellulose, e.g. carboxy-methyl, -ethyl or -propylcellulose. These are often present, when used e.g. in agrochemical formulations in particular as 0.01 to 5 wt% of a concentrate formulation and correspondingly diluted in a spray formulation at end use dilution.

The agrochemical compositions of the invention can further include a chemical agent which makes the activity of the agrochemical more specific to or selective in the intended use of the composition. The chemical agent can be a phytotoxicity inhibitor, to regulate any herbicidal activity of the concentrate or corresponding dilute formulation, or to make it more selective, e.g. as between graminious and broad-leaved species. Such a second chemical agent may be present as up to 10 wt% of concentrated compositions and typically up to 0.25% of diluted compositions for spraying.

The compositions of the invention can be made relatively straightforwardly, usually by simple mixing of the components. Conveniently, this may be carried out by dissolving the active agrochemical(s) and the surfactants and any other components in water to give either a concentrate for subsequent dilution to end use concentrations or directly at end use concentration e.g. in the spray tank.

Concentrated agrochemical compositions can be converted into spray formulations by dilution with water, typically in an amount of from 10 to 10,000 times the total weight of the agrochemical and adjuvant components e.g. with 30 to 1,000 times the total weight of these components.

- 11 -

The following Examples illustrate the invention. All parts and percentages are by weight unless otherwise stated.

Materials

AS1	aqueous C ₈ -C ₁₁ alkyl saccharide (DP between 1.6 and 1.9) adjuvant ex ICI (about 70% alkyl saccharide)
AS2	aqueous C ₈ -C ₁₁ alkyl saccharide (DP between 1.2 and 1.5) adjuvant ex ICI (about 50% alkyl saccharide)
PEG 300	polyethylene glycol average molecular weight about 300 from ICI
PEG 400	polyethylene glycol average molecular weight about 400 from ICI
Touchdown	Proprietary glyphosate trimethylsulphonium salt herbicide formulation including alkyl saccharide adjuvant ex Zeneca
ETA	conventional ethoxylated tallow amine adjuvant (95+% active)
water	the water used to make up the spray formulations used in the Examples had a standard hardness of 342 ppm.

Methods

Test spray formulations of sulfosate herbicide were made by dissolving the various components in water. The sulfosate concentration used was 330 g.l⁻¹ and the adjuvant compositions 240 g.l⁻¹ of adjuvant (alkyl glycoside including PEG). These compositions were diluted to a concentration of 4.95 g.l⁻¹ sulfosate (1.5 litres of concentrates composition per 100 litres of spray). Weed control using glyphosate type herbicides was assessed by spraying the diluted herbicide test formulations onto 6 m x 2.5 m in 4 replicate sets of test plots in a fallow field. The spray volume used was 200 l.ha⁻¹ corresponding to 3 l.ha⁻¹ of concentrate and 990 g.ha⁻¹ of active sulfosate. The field flora were predominantly grasses with a roughly even spread of some perennial dicotyledon weeds. At the time of application of the test sprays, the field flora included *Ranunculus repens* (flowering 15 to 30 cm high), *Rumex obtusifolius* (in seed 30 to 50 cm), *Taraxacum officinale* (15 to 20 cm flowering), *Trifolium repens* (15 to 25 cm), *Cirsium arvense* (20 to 25 cm), *Urtica dioica* (30 to 50 cm) *Holcus lanatus* (flowering 30 to 40 cm) and *Poa trivialis* (15 to 20 cm). The weather at application and throughout the test period was dry and sunny and this may have affected the results.

Herbicidal activity was assessed by visual estimation of the percentage remaining green foliage after 15 and 28 days after spraying in each test plot compared to the control plots. Estimates of the

- 12 -

control of growth of the specific species *Ranunculus repens* (RANRE) and *Rumex obtusifolius* (RUMOSS) in treated pots as compared with the control plots was also estimated at 15 days after spraying. These data were scored as a percentage.

Comparison formulations included glyphosate type herbicide without any adjuvant, glyphosate type herbicide with alkyl glucoside as adjuvant (without any PEG) and commercial formulations of glyphosate type herbicide (including any built in adjuvants).

Examples

Various herbicide formulations were made up as set out in Table 1 below.

Table 1

Ex No	Alkyl saccharide		PEG		AS:PEG ratio	Other Surfactant		Application rate		
	nature	(%)	nature	(%)		nature	amount	(l.ha ⁻¹)	(g.ha ⁻¹)	
C1	no treatment									
C2	Touchdown							3	990	
C3		100	-	0	-	-	-	3	990	
1	AS1	75	PEG 300	25	2.1	-	-	3	990	
2	AS1	75	PEG 300	20	2.8	ETA	5	3	990	
3	AS2	75	PEG 300	25	2.1	-	-	3	990	

The results of weed control trial using the protocol described above are set out in Table 2 below:

Table 2

Ex No	% green weed after:		% control of species:	
	2 weeks	4 weeks	RANRE	RUMOSS
C1	100	100	0	0
C2	19	17.5	83	27.5
C3	22.5	22.5	85	32.5
1	16	26	83	22.5
2	21	27.5	87	37.5
3	19	20	78	50

Claims

- 1 A composition which comprises
 - i at least one hydrocarbyl polysaccharide;
 - ii at least one polyalkylene glycol; and
 - iii water.
- 2 A composition as claimed in claim 1 in the form of a water dispersible and/or soluble agrochemical composition comprising:
 - i at least one hydrocarbyl polysaccharide;
 - ii at least one polyalkylene glycol;
 - iii water; and
 - iv at least one agrochemical.
- 3 A composition as claimed in claim 2 in the form of an agrochemical composition which is dispersible in water and is itself a stable homogeneous aqueous solution comprising:
 - i at least one hydrocarbyl polysaccharide;
 - ii at least one polyalkylene glycol;
 - iii water; and
 - v at least one glyphosate type herbicide.
- 4 A composition as claimed in any one of claims 1 to 3 which includes at least one other water soluble or water dispersible non-ionic surfactant.
- 5 A composition as claimed in any one of claims 1 to 4 wherein the hydrocarbyl polysaccharide is at least one at least one compound of the formula:

$$\text{ROG}_a \quad \text{where}$$
 - R is a hydrocarbyl group;
 - G is a saccharide residue; and
 - a is an average value which is at least 1.
- 6 A composition as claimed in claim 5 wherein the hydrocarbyl polysaccharide is at least one glucoside compound of the formula:

$$\text{R}'\text{O}(\text{C}_6\text{H}_{10}\text{O}_5)_a\text{H} \quad \text{where}$$

- 14 -

R' is a C₈ to C₁₄ alkyl group; and

a' is between 1 and 2.

- 7 A composition as claimed in any one of claims 1 to 6 wherein the polyalkylene glycol is or includes at least one polyethylene glycol, polypropylene glycol or co-poly(ethylene/propylene) glycol.
- 8 A composition as claimed in claim 7 wherein the polyalkylene glycol is or includes polyethylene glycol having a molecular weight of from about 200 up to about 2000.
- 9 A composition as claimed in claim 8 wherein the polyethylene glycol has a molecular weight of from about 200 to about 500.
- 10 A composition as claimed in any one of claims 1 to 9 wherein the weight ratio of hydrocarbyl saccharide to polyalkylene glycol is in the range of about 20:1 to 1:2.
- 11 A composition as claimed in claim 10 wherein the polyalkylene glycol is polyethylene glycol and the weight ratio of hydrocarbyl saccharide to polyethylene glycol is about 10:1 to about 1.5:1.
- 12 A composition as claimed in any one of claims 1 to 11 wherein the composition includes at least one other non-ionic surfactant of the formulae:

$$R^1 O(X)R^2 \quad \text{and/or} \quad R^3 - N(-XH)_2 \quad \text{in which}$$

R¹ is a C₈ to C₁₈ alkyl, alkyl phenyl, a sorbitan or a C₁₀ to C₂₂ fatty acyl group;

R² is hydrogen or an alkyl or carboxyalkyl group or a salt thereof, or a C₁₀ to C₂₂ fatty acyl group;

X is a polyalkylene oxide group having an average of 2 to 40 alkylene oxide groups or mixed polyalkylene oxide groups, and

R³ is a C₁₀ to C₂₂ alkyl or alkenyl group.
- 13 A composition as claimed in claim 12 wherein the amount of the other non-ionic surfactant is from 5% to 120% by weight of the alkyl glycoside.

- 15 -

- 14 A composition as claimed in any one of claims 1 to 13 which is an agrochemical adjuvant formulation wherein the proportions of the components are:
- i hydrocarbyl polysaccharide 25 to 70%;
 - ii at least one polyalkylene glycol 2 to 35%; and
 - iii water 15 to 70%;
 - iv other non-ionic surfactant (when present) 2 to 35; and
 - v other (minor) components 0 to 5%;
- the percentages being by weight based on the total of components i to v.
- 15 A composition as claimed in any one of claims 1 to 13 formulated as a dilutable agrochemical concentrate and which contains:
- a agrochemical at a concentration of from 100 to 500 g.l⁻¹;
 - b adjuvant, being the combination of hydrocarbyl polysaccharide, polyalkylene glycol and other non-ionic surfactant (when present), 30 to 500 g.l⁻¹;
 - c minor components 0 to 50 g.l⁻¹; and
 - d the remainder as water.
- 16 A method of killing or inhibiting vegetation by applying to the vegetation or the soil near or surrounding the vegetation an agrochemical composition including one or more growth regulators and/or herbicides in combination with a composition as claimed in any one of claims 1 to 15.
- 17 A method as claimed in claim 16 wherein the agrochemical is or includes at least one glyphosate type herbicide.
- 18 A method of killing or inhibiting plant pests by applying to vegetation or the soil surrounding the vegetation an agrochemical composition including one or more pesticides in combination with a composition as claimed in any one of claims 1 to 15.
- 19 A method as claimed in claim 18 wherein the agrochemical is or includes at least one insecticide, fungicide and/or acaricide.
- 20 A method as claimed in any one of claims 16 to 19 wherein the agrochemical composition is sprayed onto the vegetation at a rate of about 100 to 400 l.ha⁻¹.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/02951

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A01N25/30 A01N57/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 00010 A (ZENECA LTD ;TOWNSON JANE KAREN (GB); HART CLIFFORD ARTHUR (GB); OS) 4 January 1996 see page 7, line 10-20; example 7; table 9 ---	1-20
P, X	WO 98 39273 A (RHODIA) 11 September 1998 see page 8, line 11-13; page 10, line 9/10; page 11, line 18 - page 12, line 21; example iii/table iii. ---	1,2,4-8, 10,14,15
X	EP 0 353 735 A (KAO CORP) 7 February 1990 see table 1 --- -/--	1,2,4-8, 10,14,15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 December 1998

Date of mailing of the international search report

30/12/1998

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Klaver, J

INTERNATIONAL SEARCH REPORT

Int. Patent Application No

PCT/GB 98/02951

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 12259 A (HOORNE DIRK ;AUDA MAHROUSSA (BE); ICI PLC (GB)) 9 June 1994 cited in the application see the whole document ---	1-20
Y	EP 0 220 902 A (STAUFFER CHEMICAL CO) 6 May 1987 cited in the application see the whole document ---	1-20
Y	EP 0 498 785 A (MONSANTO EUROPE SA) 12 August 1992 see page 4, line 12-14; example 5 ---	1-20
Y	DE 36 36 994 A (STAUFFER CHEMICAL CO) 7 May 1987 see page 3, line 42-64; tables III, IV ---	1-20
Y	EP 0 526 444 A (MONSANTO CO) 3 February 1993 see page 4, line 6-25 ---	1-20
Y	EP 0 364 202 A (ICI AUSTRALIA LTD) 18 April 1990 see page 2, line 31-42 -----	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02951

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9600010	A	04-01-1996	AU 698255 B	29-10-1998
			AU 3966195 A	19-01-1996
			BR 9508055 A	12-08-1997
			CA 2189880 A	04-01-1996
			CN 1151103 A	04-06-1997
			EP 0767609 A	16-04-1997
			EP 0852113 A	08-07-1998
			JP 10502071 T	24-02-1998
			NO 965550 A	21-02-1997
			PL 317943 A	12-05-1997
			ZA 9504749 A	13-05-1996
WO 9839273	A	11-09-1998	NONE	
EP 0353735	A	07-02-1990	JP 2043294 A	13-02-1990
			JP 2582626 B	19-02-1997
			JP 2060999 A	01-03-1990
			JP 2566816 B	25-12-1996
			JP 2069597 A	08-03-1990
			JP 2526102 B	21-08-1996
			JP 2123195 A	10-05-1990
			JP 2530215 B	04-09-1996
			DE 68925390 D	22-02-1996
			DE 68925390 T	05-06-1996
			ES 2084591 T	16-05-1996
			HK 108096 A	05-07-1996
			US 5230835 A	27-07-1993
WO 9412259	A	09-06-1994	US 5750513 A	12-05-1998
			AU 676518 B	13-03-1997
			AU 5469294 A	22-06-1994
			EP 0671967 A	20-09-1995
			JP 8503882 T	30-04-1996
			NZ 257878 A	29-01-1997
			SG 46274 A	20-02-1998
			ZA 9308590 A	01-06-1994
EP 0220902	A	06-05-1987	AT 164726 T	15-04-1998
			AU 591938 B	21-12-1989
			AU 6419186 A	30-04-1987
			BR 8605102 A	21-07-1987
			DE 3650670 D	14-05-1998
			DE 3650670 T	30-07-1998
			DK 504886 A	22-04-1987
			ES 2113844 T	16-05-1998
			FI 864231 A, B,	22-04-1987
			GR 3026574 T	31-07-1998
			JP 2050658 C	10-05-1996
			JP 7084367 B	13-09-1995
			JP 62099313 A	08-05-1987
			US 5468718 A	21-11-1995
EP 0498785	A	12-08-1992	AT 161144 T	15-01-1998
			AU 650091 B	09-06-1994
			AU 1060092 A	13-08-1992
			CA 2060870 A	09-08-1992
			DE 69223529 D	29-01-1998
			DE 69223529 T	02-07-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02951

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0498785 A		DK 498785 T	09-02-1998
		ES 2112308 T	01-04-1998
		GR 3026188 T	29-05-1998
		IL 100873 A	18-02-1997
		JP 4352704 A	07-12-1992
		KR 9509173 B	16-08-1995
		NZ 241500 A	26-10-1994
		PT 100103 A	31-05-1993
DE 3636994 A	07-05-1987	AU 586293 B	06-07-1989
		AU 6455286 A	07-05-1987
		BR 8605400 A	11-08-1987
		FR 2589328 A	07-05-1987
		JP 62120304 A	01-06-1987
EP 0526444 A	03-02-1993	US 5430005 A	04-07-1995
		AT 146035 T	15-12-1996
		AU 648919 B	05-05-1994
		AU 2066192 A	04-02-1993
		CA 2075004 A	03-02-1993
		DE 69215785 D	23-01-1997
		DE 69215785 T	26-06-1997
		JP 6024917 A	01-02-1994
		MX 9204470 A	01-05-1993
		NZ 243780 A	23-12-1993
		ZA 9205731 A	05-11-1993
EP 0364202 A	18-04-1990	AT 126667 T	15-09-1995
		AU 627503 B	27-08-1992
		AU 4243489 A	26-04-1990
		CA 2000513 A	13-04-1990
		DE 68923926 D	28-09-1995
		DE 68923926 T	22-02-1996
		GR 3017209 T	30-11-1995
		JP 2169504 A	29-06-1990
		JP 2726715 B	11-03-1998